The Supersymmetric Higgs

Puneet Batra & Eduardo Ponton Columbia University arXiv:0809.3453 [hep-ph]

The MSSM Scalar Sector

First choice for non-SM Higgs searches

2 parameters ———— Physics of 4 fields
$$(m_A, t_\beta)$$
 (h^0, H^0, H^\pm, A^0)

a generic 2HDM has 7 more parameters...

The MSSM Scalar Sector


First choice for non-SM Higgs searches

2 parameters ———— Physics of 4 fields
$$(m_A, t_\beta)$$
 (h^0, H^0, H^\pm, A^0)

a generic 2HDM has 7 more parameters...

Radiative EWSB (electroweak symmetry breaking)

Radiative EWSB (electroweak symmetry breaking)




$$Tr = -\frac{1}{2}M_Z^2 - \frac{2\sec^2(2\beta)}{M_Z^2}Det$$



No EWSB w/o SUSY breaking

 $Det(M^2)$

Radiative EWSB (electroweak symmetry breaking)



$$Tr = -\frac{1}{2}M_Z^2 - \frac{2\sec^2(2\beta)}{M_Z^2}Det$$



No EWSB w/o SUSY breaking

$$Det(M^2)$$

 $m_{H_u}^2$ driven negative by top loops

Beyond The MSSM Scalar Sector

Very Light SM-like Higgs mass

$$m_{h^0} < M_Z$$
 at tree-level $m_{h^0} \leq 130~{
m GeV}$ with %-ish tuning

Beyond The MSSM Scalar Sector

Very Light SM-like Higgs mass

$$m_{h^0} < M_Z$$
 at tree-level $m_{h^0} \leq 130~{
m GeV}$ with %-ish tuning

• Can use strong-coupling to increase $\,m_h$ o

```
m_{h^0} \sim 300~{
m GeV} Haber-Sher, Espinosa-Quiros, ... P.B. et al., "The Fat Higgs", ..., Dine et al.
```

Beyond The MSSM Scalar Sector

Very Light SM-like Higgs mass

$$m_{h^0} < M_Z$$
 at tree-level $m_{h^0} \leq 130~{
m GeV}$ with %-ish tuning

• Can use strong-coupling to increase $\,m_h^{_0}$

$$m_{h^0} \sim 300~{
m GeV}$$
 Haber-Sher, Espinosa-Quiros, ... P.B. et al., "The Fat Higgs", ..., Dine et al.

Can evade LEP bounds via singlet mixing

NMSSM, Dermisek-Gunion, Chang-Fox-Weiner

$$h^0 \rightarrow 2a \rightarrow 4b, 4\tau$$

big modifications in the higgs sector! affects collider searches

Outline

• LEP motivates Supersymmetric EWSB (sEWSB)

SM-like Higgs mass is not directly tied to M_Z Another qualitative shift in Higgs physics

sEWSB captured in an effective field theory

Only need to consider the MSSM degrees of freedom Much easier to identify gross phenomenology (Fat Higgs)

New paradigm for 2HDM phenomenology

The heavier higgs, H^0 , is naturally SM-like The charged Higgs and non-SM-like Higgs are degenerate Very rich vacuum structure

Supersymmetric EWSB (sEWSB)

SUSY-limit of EWSB

Vector Superfield 'eats' a chiral superfield

$$M_{H^{\pm}} = M_{W^{\pm}}$$
 , $M_{h^0} = M_{Z^0}$


The super-radial mode contains the SM-like Higgs

$$\rho\supset (H^0,A^0)$$


• $m_{
ho}$ determined by the superpotential W, not g_w , much easier to find $m_{
ho}>114~{
m GeV}$

$$W \supset \mu H_u H_d + \frac{\omega_1}{2\mu_S} (H_u H_d)^2$$

$$W \supset \mu H_u H_d + \frac{\omega_1}{2\mu_S} (H_u H_d)^2$$



$$W \supset \mu H_u H_d + \frac{\omega_1}{2\mu_S} (H_u H_d)^2$$



The EFT has a good expansion parameter

Corrections from higher-order operators

$$\sim \frac{v^2}{\mu_S^2} = \frac{1}{\omega} \frac{\mu}{\mu_S}$$

The EFT has a good expansion parameter

Corrections from higher-order operators

$$\sim \frac{v^2}{\mu_S^2} = \frac{1}{\omega} \frac{\mu}{\mu_S}$$

Irony: under control only because there is no renormalizable tree-level quartic--- the usual problem in the MSSM!

The EFT has a good expansion parameter

Corrections from higher-order operators

$$\sim \frac{v^2}{\mu_S^2} = \frac{1}{\omega} \frac{\mu}{\mu_S}$$

Irony: under control only because there is no renormalizable tree-level quartic--- the usual problem in the MSSM!

• Some tension between making m_{ρ} large, but keeping the EFT under control.

Inverted scalar hierarchy occurs when $~2\mu > M_Z$


Turn on SUSY-breaking

SUSY-breaking needed to lift the fermions

Mass	Scalars	Fermions	Vectors
0		1 majorana	A_{μ}
m_W	H^\pm	2 Dirac	W_{μ}^{\pm}
m_Z	H	1 Dirac	$Z_{\mu}^{'}$
$2 \mu $	h, A^0	1 majorana	<u> </u>

- Breaks the degeneracy with the origin
- Many terms, even without any new degrees of freedom

Much larger region of EWSB



Generically, much larger region for EWSB


signs matter,


$$eta \in \left[-rac{\pi}{2}, rac{\pi}{2}
ight]$$


$$\tan(\beta) < 0!$$

Inverted Hierarchy in the MSSM

• Inverted Hierarchy: H^0 is SM-like ($m_{h^0} < m_{H^0}$)








$$M_{\tilde{t}} \sim m_t$$


 $M_{\tilde{t}} \sim 400 \; \mathrm{GeV}$

$$M_{\tilde{t}} \sim 600 \text{ GeV}$$

Must satisfy $m_A^2 \sim m_d^2 - m_u^2 - m_Z^2$ to a high degree

Inverted Hierarchy in sEWSB vacua





Inverted hierarchy in roughly 'half' of parameter space

Example Spectra

Inverted Spectra are easy to find:

μ	ω	μ/μ_s	b/μ^2	m_u^2/μ^2	$m_{H_d}^2/\mu^2$	ξ	M_1/μ	M_2/μ
-150	2	0.14	-1.1	-0.99	-0.51	0.20	0.36	0.57

ρ	$\tan \beta$	m_{h^0}	m_{H^0}	$g_{H^0ZZ}^2/g_{h_{\rm SM}ZZ}^2$	m_{A^0}	m_{H^+}	m_{χ^+}	$oxed{m_{\chi^0}}$
	-1.3					190	l	I I

$$H^+ \rightarrow \chi^+ \chi^0$$

Chargino NLSP?

μ	ω	μ/μ_s	b/μ^2	m_u^2/μ^2	$m_{H_d}^2/\mu^2$	ξ	M_1/μ	M_2/μ
-70	1	0.11	-1.6	-1.7	.22	0.20	1.5	1.7

ρ	$\tan \beta$	m_{h^0}	m_{H^0}	$g_{H^0ZZ}^2/g_{h_{\rm SM}ZZ}^2$	m_{A^0}	m_{H^+}	m_{χ^+}	$oxed{m_{\chi^0}}$
0.34	-1.8	120	140	0.82	110	125	100	110

Conclusions

 Post-LEP, it is worth reconsidering what the most likely BSM Higgs sector looks like:

$$h^0
ightarrow 2a
ightarrow 4b$$
 $m_{h^0} < m_{H^0}, H^0$ SM-like

 Supersymmetric EWSB is a qualitatively new starting point---EFT approach is very powerful!

Easy to UV complete into a theory with $~W \supset \lambda S H_u H_d$

• Light Higgs -> Light charginos, charged Higgs + new phenomenology!

Rich Vacuum Structure---cosmological applications?

• supplements

Decoupling of sEWSB vacuum

